

60 Years of the Invertebrate Primitive Antibody (IPA): A Paradigm Shift in Evolutionary Immunology

Michel Leclerc*

Immunology of Invertebrates, Department of Biochemistry and Biology, Orléans University, Orléans, France.

***Correspondence:**

Michel Leclerc, Immunology of Invertebrates, Department of Biochemistry and Biology, Orléans University, Orléans, France.

Received: 01 Sep 2026; **Accepted:** 02 Oct 2026; **Published:** 11 Oct 2026

Citation: Michel Leclerc. 60 Years of the Invertebrate Primitive Antibody (IPA): A Paradigm Shift in Evolutionary Immunology. Japanese J Med Res. 2026; 4(4): 1-3.

ABSTRACT

For over six decades, the mainstream consensus in evolutionary immunology has maintained the central dogma that adaptive immunity—defined by clonal expansion, specific immunoglobulins, and dedicated lymphocytes—is an exclusive innovation of jawed vertebrates. This review synthesizes sixty years of structural, functional, and genomic evidence demonstrating that the sea star (Echinodermata) possesses a fully operational, primitive adaptive immune system. We detail the anatomical characterization of sea star B and T lymphocytes within ancestral lymphoid organs, trace the functional mechanics of the Invertebrate Primitive Antibody (IPA), and present modern genomic data confirming the existence of MHC-like networks. By systematically addressing the structural homologies between echinoderm and mammalian immune cells, this paper directly challenges conventional evolutionary dogmas and proposes a unified framework for the origin of adaptive immunity.

Keywords

Invertebrate Primitive Antibody (IPA), B sea star Lymphocyte, T sea star Lymphocyte, Adaptive Immunity, Echinoderm Genomics, Axial Organ.

Introduction: Deconstructing the Vertebrate Dogma

The historical narrative championed by mainstream bodies, such as the American Association of Immunologists (AAI), dictates that invertebrates rely strictly on non-specific, germline-encoded innate defense mechanisms. This paradigm asserts that in the absence of "true" lymphocytes and functional antibodies, clonal expansion is impossible.

However, sixty years of continuous empirical investigation into echinoderm biology severely undermines this assumption. Rather than an abrupt evolutionary leap occurring exclusively at the vertebrate boundary, experimental evidence points to a deep, invertebrate origin of specific immunity. This review contextualizes these findings across three fundamental aspects: cellular morphology, adaptive functionality mediated by the Invertebrate Primitive Antibody (IPA), and comparative genomics [1].

Cellular Architecture: Sea Star B and T Lymphocytes

The Sea Star B Lymphocyte

The sea star B lymphocyte(2) is primarily localized within the axial organ—an asymmetrical, specialized structure functioning as an ancestral lymphoid organ—as well as Tiedemann's bodies across various starfish species.

- **Ultrastructural Characteristics:** Transmission Electron Microscopy (TEM) reveals a cell approximately 4 μm in diameter featuring a highly developed, prominent rough endoplasmic reticulum (RER). The overall cytomorphology mirrors that of mammalian plasma cells actively engaged in protein synthesis.
- **Surface Receptor Homologies:** Phenotypic analysis via flow cytometry establishes that these cells actively bind human T-Cell Receptor (TCR) analogues and the CD14 human receptor.
- **Clonal Proliferation:** Following in vivo immunization with diverse foreign antigens, the sea star B lymphocyte population undergoes massive expansion within the axial organ, shifting from a baseline of 10% to upwards of 45% of the total cellular population.

The Sea Star T Lymphocyte

Isolated alongside B cells in the axial organ, the sea star T lymphocyte [2] represents the cellular arm of echinoderm adaptive defense.

- **Biophysical and Mitogenic Properties:** These 4 μm diameter cells exhibit distinct physical properties, including non-adherence to nylon wool. They undergo selective agglutination and mitogenic stimulation when exposed to lectins such as Concanavalin A (ConA) and Pokeweed Mitogen (PWM), matching the classic activation pathways of mammalian T cells.
- **Functional Assays:** When transplanted into vertebrate hosts, sea star T lymphocytes successfully induce physiological hallmarks of cell-mediated immunity, including angiogenesis and splenomegaly.
- **Receptor Affinity:** Flow cytometry confirms that sea star T lymphocytes possess binding affinities for CD7 human receptors and TCR profiles (though the latter is expressed at a lower relative degree than seen in B-like counterparts).

The Invertebrate Primitive Antibody (IPA) and Adaptive Mechanics

The functional execution of sea star adaptive immunity relies on a sophisticated tripartite cellular cooperation first elucidated experimentally in 1986. The process mimics the classical vertebrate immune response:

1. **Antigen Processing:** Non-specific phagocytes isolate and process foreign antigens, effectively serving the functional role of primitive macrophages.
2. **Cellular Cooperation:** Processed antigens facilitate direct cooperative interactions between sea star T lymphocytes and sea star B lymphocytes.
3. **Humoral Secretion:** This cellular cascade culminates in the synthesis and secretion of the Invertebrate Primitive Antibody (IPA), generating highly specific, inducible immune responses capable of targeted hapten and protein neutralization [4].

Echinoderm Genomics and the Ig-Kappa-like / MHC Framework [3]

The structural and cellular assertions of sea star adaptive immunity are directly supported by recent high-throughput genomic and transcriptomic sequencing. Decisive genomic profiling has successfully identified the molecular blueprints underpinning the Invertebrate Primitive Antibody (IPA) and self/non-self recognition networks:

Structural Integrity of the Sea Star Ig-kappa Gene [5]

Molecular cloning and transcriptomic mapping of the sea star genome have confirmed the presence of explicit Immunoglobulin kappa (Ig-kappa) light-chain like sequences.

- **Variable (V) and Constant (C) Domain Conservation:** Genomic analysis reveals that the sea star Ig-kappa-like gene maintains highly conserved regions homologous to vertebrate V and C domains. The primary sequence preserves the definitive cysteine residues required to form the classical intrachain disulfide bonds, stabilizing the structural immunoglobulin fold.
- **Primitive V(D)J Geometry:** Unlike the hyper-diversified somatic recombination found in higher mammals, the echinoderm Ig-kappa architecture represents a critical evolutionary intermediate—demonstrating that the genomic capability to express light-chain analogues predates the separation of proto-chordates and echinoderms.

The MHC Core Matrix

Complementing the humoral antibody-like network, genomic sequencing has successfully mapped:

- **MHC Class Gene Homologies:** Conserved sequences encoding primitive Major Histocompatibility Complex (MHC) gene orthologues. These structures provide the mandatory genetic apparatus required for allorecognition, tissue graft rejection, and specific antigen presentation to T-like cells.

Comparative Immunology Synthesis

To illustrate the functional and structural evolutionary bridge between the Echinodermata adaptive system and modern mammals, the key immunological traits established over 60 years of empirical research are synthesized below:

Conclusion: A Direct Reply to Evolutionary Immunologists

The persistence of the term "true lymphocyte" in contemporary literature functions more as a semantic barrier than a biological reality. If a cell possesses identical ultrastructural morphology, undergoes targeted clonal expansion upon immunization, displays mitogenic responses to specific lectins, carries homologous surface clusters of differentiation, and secretes specific neutralizing factors, it is a lymphocyte.

Sixty years of IPA research proves that the molecular machinery of adaptive immunity did not appear spontaneously in jawed

Immunological Trait	Sea Star System (Echinodermata)	Mammalian System (Vertebrata)
Ancestral Lymphoid Organ	Axial organ / Tiedemann's bodies	Lymph nodes, spleen, thymus
B-Cell Functional Homologue	B sea star lymphocyte (4 μm, high RER density)	B lymphocyte / Plasma cell
T-Cell Functional Homologue	T sea star lymphocyte (4 μm, nylon-wool non-adherent)	T lymphocyte
Mitogenic Inducibility	ConA / Pokeweed Mitogen (PWM) response	Lectin-induced blastogenesis
In Vivo Cellular Proliferation	Antigen-induced clonal shift (10% up to 45%)	Clonal expansion in germinal centers
Humoral Mediator	Invertebrate Primitive Antibody (IPA)	Immunoglobulins (IgG, IgM, IgA, etc.)
Genomic Light-Chain Orthologue	Ig-kappa-like gene (Conserved V/C domains)	Ig-kappa light chain gene
Allorecognition Matrix	Primitive MHC class genes	Classical MHC Class I and Class II complexes

vertebrates. Instead, the evolutionary foundation was laid millions of years prior within the ancestral lymphoid organs of the Echinodermata.

References

1. Leclerc M, Brillouet C, Luquet G. The axial organ of the sea star *Asterias rubens*: An ancestral lymphoid organ. *Dev Comp Immunol*. 1980; 4: 605-615.
2. Leclerc M, Arneodo VJ, Legac E, et al. Identification of T-like and B-like lymphocyte subsets in sea star *Asterias rubens* by monoclonal antibodies to human leukocytes. *Thymus*. 1993; 21: 133-139.
3. Leclerc M, Loïc Baerlocher. Mapping on sea star MHC genes in Invertebrates. *European Journal of Biology and Biotechnology*. 2021; 2: 60-62.
4. Leclerc M, Brillouet C, Luquet G, et al. Production of an antibody-like factor in the sea star *Asterias rubens*: involvement of at least three cellular populations. *Immunology*. 1986; 57: 479-482.
5. Nadine Vincent, Magne Osteras, Patricia Otten, et al. A new gene in *Asterias rubens*: A sea star IGHKappa gene. *Meta Gene*. 2014; 2: 320-322.