

Benign Squamous Papilloma of the Vulva

Sowjanya Kurakula¹, Vandana Muralidharan², Shravani Rama³, Abhijna Rao Kompella⁴, Navya N⁵

¹Department of Obstetrics and Gynecology, Former Head, Sekgoma Memorial Hospital, Serowe, Botswana, Africa.

²Department of Obstetrics and Gynecology, Lifeline Medical Associates, New Jersey, USA.

³Department of Obstetrics and Gynecology, Gandhi Medical College, Secunderabad, India.

⁴Department of Obstetrics and Gynecology, Srinivasa Maternity and Nursing Home, Hyderabad, India.

⁵Department of Obstetrics and Gynecology, Ranibow Children's Hospital and BirthRight, Bengaluru, India.

*Correspondence:

Sowjanya Kurakula, Department of Obstetrics and Gynecology, Former Head, Sekgoma Memorial Hospital, Serowe, Botswana, Africa.

Received: 12 Apr 2023; Accepted: 16 May 2023; Published: 20 May 2023

Citation: Kurakula S, Muralidharan V, Rama S, et al. Benign Squamous Papilloma of the Vulva. Gynecol Reprod Health. 2023; 7(3): 1-1.



The benign squamous papilloma of the vulva is a rare skin condition. It is often considered a skin tag variant. Our case is an 18-year-old female who presented to the outpatient department with a slow-growing painless vulvar growth for the past 6 years. It was located on labium majus, 10×7cm in size, soft, mobile, non-tender, pedunculated, and irregular in shape and growth. Her basic investigations are Hemoglobin 11.2 gm%, hematocrit 33%, MCV 80 μm³, urinalysis within normal limits, HIV & HbsAg negative. An excision biopsy was done for esthetic reasons. The histopathology report revealed acanthotic epithelium, hyperkeratosis, and papillary projections

with the fibrovascular core. Her postoperative recovery was uneventful. She presented at one monthly follow-up with sound healing. We bring forward this case as it is one of the uncommon benign masses of the vulva.

Final Diagnosis

Benign squamous papilloma of the vulva.

Three Differential Diagnosis

Achrocordon, Genital warts, Lipoma.