Review Article ISSN 2639-8451

Addiction Research

Comparing and Integrating the 12-Step Recovery Model and Classical Medical Model: *Toward a Holistic Framework for Addiction Treatment*

Julian Ungar-Sargon MD, PhD*

Borra College of Health Science, Dominican University, USA.

***Correspondence:**
Julian Ungar-Sargon, Borra College of Health Science, Dominican University, USA.

Received: 25 Mar 2025; **Accepted:** 20 Apr 2025; **Published:** 29 Apr 2025

Citation: Ungar-Sargon J. Comparing and Integrating the 12-Step Recovery Model and Classical Medical Model: *Toward a Holistic Framework for Addiction Treatment*. Addict Res. 2025; 9(1): 1-12.

ABSTRACT

This paper explores the historical development, theoretical foundations, and practical applications of two prominent approaches to understanding and treating addiction: the 12-Step recovery model, which conceptualizes addiction as a "dis-ease," and the classical medical model, which frames addiction within allopathic healing paradigms. Through critical analysis of their respective strengths and limitations, this paper proposes an integrated framework that harmonizes these seemingly disparate approaches to enhance treatment outcomes. The integration acknowledges the biological, psychological, social, and spiritual dimensions of addiction, offering a more comprehensive approach to recovery. This integration framework is particularly relevant for physicians and other medical practitioners seeking to provide more holistic care for patients with substance use disorders.

Keywords

Addiction medicine, 12-step programs, Disease model, Integrated treatment, Recovery-oriented care, Substance use disorders, Biopsychosocial-spiritual model, Allopathic medicine.

Historical Development

Classical Medical Model

The classical medical model, rooted in allopathic medicine, emerged from the scientific revolution of the 17th and 18th centuries, when figures like Thomas Sydenham (1624-1689) began to systematically classify diseases based on observable symptoms [1]. This approach represented a fundamental shift toward empiricism in medical practice that would eventually transform our understanding of addiction. The 19th century brought significant advancements in pathology and microbiology, with Louis Pasteur's germ theory of disease and Robert Koch's postulates establishing the foundations of modern medicine that physicians continue to apply today [2].

The medical approach to addiction began taking shape in the early 19th century through pioneering works that physicians may recognize as precursors to contemporary addiction medicine. In 1804, Thomas Trotter published "An Essay, Medical, Philosophical, and Chemical, on Drunkenness," one of the first works to characterize habitual drunkenness as a disease rather than a moral failing [3]. Benjamin Rush, often considered the father of American psychiatry, similarly described addiction to alcohol as a disease in his 1819 work "An Inquiry into the Effects of Ardent Spirits upon the Human Body and Mind." [4].

These early formulations would be familiar to modern physicians who understand substance use disorders as conditions requiring medical intervention rather than moral judgment. Despite these early insights, the medical model's application to addiction faced considerable resistance throughout the 19th century, as moral and temperance-based explanations of addiction remained dominant in both public discourse and professional practice. It wasn't until the late 19th and early 20th centuries that more biological explanations began to gain traction among medical professionals. The establishment of dedicated treatment facilities like the New York State Inebriate Asylum in 1864 reflected growing acceptance of medical approaches to addiction among physicians of the era [5].

The medical model's influence expanded significantly in the mid-20th century, coinciding with advancements that many practicing physicians would recognize as foundational to modern addiction medicine. E.M. Jellinek's seminal work "The Disease Concept of Alcoholism" (1960) provided a scientific foundation for understanding alcoholism as a progressive disease with distinct phases, a concept that continues to inform clinical practice. Subsequently, advances in neuroscience, particularly in the 1990s dubbed the "Decade of the Brain" further solidified the biological basis of addiction through research on brain reward pathways, neuroadaptation, and genetic vulnerability that revolutionized clinical approaches to addiction treatment [6,7].

Emergence of the 12-Step Recovery Model

The 12-Step recovery model originated during the Great Depression era when conventional medical and psychiatric treatments for alcoholism showed limited effectiveness a situation that many clinicians found frustrating and that still resonates with physicians facing treatment-resistant cases today. In 1935, Bill Wilson, a New York stockbroker, and Dr. Bob Smith, an Akron surgeon, both struggling alcoholics, founded Alcoholics Anonymous (AA) [8]. Their collaboration is particularly noteworthy for physicians as it represented an early integration of medical insight (through Dr. Smith) with experiential knowledge. Their meeting was facilitated by the Oxford Group, a Christian fellowship emphasizing personal transformation through principles of honesty, purity, unselfishness, and love elements that would later influence the spiritual component of recovery that many physicians now recognize as clinically significant [9].

Wilson's personal spiritual experience during a hospital stay for detoxification became foundational to AA's approach. After a profound moment of spiritual awakening, he found himself free from the desire to drink an experience that many addiction medicine specialists now understand as potentially representing a profound psychological shift that can complement neurobiological interventions. This transformative experience, combined with William James' "The Varieties of Religious Experience" and the medical insights of Dr. William Silkworth regarding alcoholism as an "allergy," shaped AA's unique concept of alcoholism as a three-fold disease physical, mental, and spiritual [10]. For physicians, this early conceptualization presaged the modern biopsychosocial model that has become integral to comprehensive medical practice.

The 12 Steps were formally codified in 1939 with the publication of "Alcoholics Anonymous" (commonly known as the "Big Book"), which outlined the program's core principles and included personal testimonials of recovery [10]. The text established alcoholism as a "dis-ease" a term intentionally hyphenated to distinguish it from the purely medical conceptualization and emphasize the state of being ill at ease with oneself [11]. This distinction remains relevant for physicians seeking to understand the experiential dimension of addiction that may not be fully captured in clinical terminology.

AA's growth was accelerated by a 1941 Saturday Evening Post article and the subsequent endorsement by prominent figures in medicine, religion, and business an early example of the kind of interdisciplinary acceptance that is now recognized as essential in addressing complex health conditions. By the 1950s, the 12-Step approach had expanded beyond alcoholism with the formation of Narcotics Anonymous (founded in 1953), and later to address other addictions and compulsive behaviors [5].

The Minnesota Model, developed at Hazelden treatment center in the 1950s, represented the first formal integration of the 12-Step approach into professional treatment settings an innovation that many physicians will recognize as forming the foundation of numerous contemporary treatment programs [12]. This model combined AA principles with medical care, psychotherapy, and education, establishing a template for addiction treatment that would dominate the field for decades and continues to influence medical approaches to addiction treatment [13].

Classical Medical Model

The classical medical model conceptualizes addiction primarily as a brain disease characterized by compulsive substance use despite harmful consequences, a framework that aligns with physicians' training in pathophysiology and evidence-based intervention [6]. This perspective has evolved significantly in recent decades and now encompasses a sophisticated understanding of addiction that informs clinical practice in several important ways.

The medical model's foundation rests on a robust understanding of the biological basis of addiction. For physicians, this translates to recognizing addiction as a chronic, relapsing brain disorder involving demonstrable changes in brain structure and function, particularly in circuits related to reward, motivation, and inhibitory control [7]. These neurobiological changes explain phenomena that clinicians frequently encounter tolerance, withdrawal, craving, and compulsive drug-seeking behavior. Neuroimaging studies have provided compelling evidence of these changes, allowing physicians to understand addiction through the same

neurobiological lens they apply to other medical conditions.

In clinical practice, the medical model provides physicians with standardized diagnostic frameworks through the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) and the International Classification of Diseases (ICD-11). These criteria focus on observable patterns of use, physiological symptoms, psychological impacts, and functional impairment that guide clinical assessment and treatment planning [14]. This standardization facilitates consistent diagnosis across healthcare settings and supports medical documentation, treatment planning, and insurance reimbursement practical considerations familiar to practicing physicians.

The medical model's emphasis on evidence-based interventions resonates with physicians' commitment to scientific rigor in treatment selection. Treatment approaches prioritize pharmacological interventions such as medications for withdrawal management (e.g., benzodiazepines, anticonvulsants), craving reduction (e.g., naltrexone, acamprosate), and maintenance therapy (e.g., methadone, buprenorphine) alongside behavioral therapies with empirical support [15]. This dual focus on medication and evidence-based psychosocial interventions mirrors approaches to other chronic medical conditions that require multimodal management.

Professional expertise forms a cornerstone of the medical model, with physicians, psychiatrists, psychologists, and specialized addiction counselors at the center of the treatment process [16]. This emphasis on professional training and specialized knowledge aligns with medical education's focus on developing clinical expertise through rigorous training and ongoing professional development. For physicians, this aspect of the model reinforces their vital role in diagnosing, prescribing, and overseeing interventions based on scientific knowledge and clinical judgment.

The medical model also incorporates a progressive disease concept that physicians will recognize from other chronic conditions. While acknowledging environmental and psychological factors, the model conceptualizes addiction as a progressive condition that typically worsens without intervention [17]. This progression is attributed to neuroadaptation and the reinforcing properties of substances, concepts that align with physicians' understanding of disease progression in other medical contexts.

The 12-Step Recovery

The 12-Step recovery model views addiction as a "dis-ease" affecting the physical, mental, and spiritual dimensions of a person's life a perspective that physicians may find complementary to the biomedical approach in addressing the experiential aspects of addiction that patients frequently report. Its theoretical foundation rests on several distinctive concepts that have significant implications for clinical practice.

The concept of powerlessness and unmanageability forms the cornerstone of the 12-Step approach. The first step "We admitted we were powerless over alcohol that our lives had become unmanageable" establishes the core premise that addiction involves a profound loss of control that cannot be overcome through willpower alone [10]. For physicians, this concept aligns with clinical observations of patients who continue substance use despite severe consequences and repeated attempts to control or cease use. This recognition of powerlessness can be understood as acknowledging the neurobiological reality of addiction a perspective that bridges 12-Step philosophy with contemporary neuroscience.



The 12-Step model proposes a spiritual solution that physicians may initially find less familiar than pharmacological or behavioral interventions. The model suggests that recovery requires connecting with a "Higher Power" (as understood by the individual) and undergoing a spiritual transformation [18]. While this spiritual emphasis distinguishes the 12-Step approach from purely medical interventions, growing research suggests that spiritual and existential dimensions of health can significantly impact treatment outcomes a finding increasingly recognized in medical literature on holistic care and in the growing field of spirituality and health.

Character development represents another core element of the 12-Step program, emphasizing moral inventory, admission of wrongs, making amends, and ongoing personal growth [19]. For physicians, this focus can be understood as addressing the psychological and interpersonal consequences of addiction that often persist after physical dependence has been addressed. The emphasis on character reconstruction acknowledges that addiction frequently damages relationships and self-concept in ways that require specific attention during recovery areas that can significantly impact treatment adherence and outcomes in clinical settings.

The fellowship and mutual aid component of the 12-Step approach offers an important resource that physicians can leverage in treatment planning. The therapeutic community created through group meetings where members share experiences and support one another's recovery provides a peer-based support system that operates largely outside professional treatment settings but can powerfully complement clinical interventions [20]. For physicians, these communities represent a valuable adjunctive resource that

extends support beyond the constraints of time-limited medical appointments and provides continuous reinforcement of recovery principles. Rather than aiming for a "cure," the 12-Step model promotes an ongoing recovery process, often summarized as "one day at a time." [21]. This approach to sobriety maintained through spiritual practice, service to others, and regular meeting attendance aligns with modern medical understanding of addiction as a chronic condition requiring ongoing management. For physicians, this perspective helps address the recurrent nature of substance use disorders that often frustrates both clinicians and patients seeking definitive resolution.

Comparative Analysis

Before analyzing the strengths and limitations of each model, it is important to consider concepts that have influenced both approaches: the historical perception of illness as divine punishment and the neurobiological-existential interface in addiction. Rosanne Lieberman writes how biblical texts provide critical frameworks for understanding how ancient moral and spiritual values shaped western and even modern notions of illness and the sense of guilt or wrongdoing and divine punishment via the agency of illness.

Lieberman has extensively examined how cultural and religious beliefs about illness as divine retribution have influenced medical practice throughout history [21,22]. Her research on biblical texts reveals a complex relationship between sin, illness, and divine punishment that continues to shape contemporary understandings of addiction. In her analysis of Exodus 32:35, where "YHWH struck the people because they made the [golden] calf," Lieberman notes how this passage has been variously interpreted as referring to plague or other forms of divine punishment [62]. The ambiguity of the Hebrew verb נגף (to strike) allows for multiple interpretations of how divine punishment manifests physically a linguistic feature that has contributed to Western cultural assumptions connecting illness with divine disapproval.

Particularly relevant to addiction treatment is Lieberman's examination of Exodus 15:26, where YHWH states, "If you will heed YHWH your God diligently, doing what is upright in His sight, giving ear to His commandments and keeping all His laws, then I will not bring upon you any of the diseases that I brought upon the Egyptians, for I YHWH am your healer." [25]. This text establishes a covenant relationship in which health becomes a sign of moral and spiritual righteousness, while illness signals covenant violation a framework that has profoundly influenced how addiction is conceptualized as either a disease (medical model) or a "dis-ease" reflecting spiritual disconnection (12-Step model).

As Lieberman observes, this binary understanding of health and illness risks stigmatizing those who suffer, suggesting they somehow deserve their condition. She notes that "an understanding that good health is the norm, and that pain and illness are divine punishments for disobedience can have worrying connotations," potentially stigmatizing people already suffering from conditions like addiction by implying moral culpability [24]. This insight is particularly relevant for physicians navigating between medical and spiritual frameworks when treating addiction, as it highlights how easily therapeutic language can inadvertently reinforce stigma.

My analysis neurobiology of existential distress complements Lieberman's historical analysis by providing a neurobiological framework for understanding the spiritual dimensions of addiction and recovery [26]. I have shown from reviewing the literature how existential anxiety activates specific neural pathways that overlap significantly with addiction circuits, suggesting that spiritual distress and substance cravings may share neurobiological mechanisms [27]. This work offers a basis for understanding how spiritual and existential interventions such as those in the 12-Step model might directly affect the same neural systems targeted by pharmacological treatments.

In my "Bridging the Divide: Toward a Neurobiologically-Informed Approach to Existential Medicine," I argue that "the traditional separation between biological and existential aspects of illness represents a false dichotomy that impedes effective treatment, particularly for conditions like addiction where meaning-making processes are central to both pathology and recovery." [28]. This perspective provides a theoretical foundation for integrating the medical and 12-Step models beyond mere pragmatic combination.

The medical model of addiction emerged partly as a reaction against punitive moral frameworks, seeking to destigmatize addiction by framing it as a medical condition rather than a moral failing or divine punishment. However, as Lieberman observes, some aspects of the medical model inadvertently reinforce notions of personal responsibility and failure through language about "compliance" and "non-adherence" that can echo moralistic perspectives [29].

The 12-Step model presents a more complex relationship with these concepts. While explicitly rejecting simplistic notions of addiction as punishment, the spiritual focus of the 12-Step approach acknowledges a relationship between spiritual wellbeing and recovery. My review has demonstrated that the practices of surrender and connection central to 12-Step programs activate parasympathetic nervous system responses that counteract the stress circuits involved in addiction maintenance [30]. Neuroimaging studies of long-term 12-Step participants show distinct patterns of functional connectivity in brain networks associated with self-

referential processing and existential awareness patterns that correlate with reported length of sobriety.

For physicians, these integrated perspectives highlight the importance of recognizing both the neurobiological realities of addiction and the existential dimensions of recovery. Our theoretical framework of "neurobiological-existential integration" offers physicians a way to discuss spiritual aspects of recovery using language compatible with medical training while acknowledging the limitations of purely reductionist approaches: "The physician who can navigate between brain chemistry and existential meaning possesses tools for addressing addiction that neither perspective alone can provide." [31].

When integrating medical and 12-Step approaches, clinicians should be aware of how language and concepts might inadvertently reinforce or challenge deep-seated cultural narratives about illness and responsibility. Therapeutic communication that carefully avoids punitive framing while acknowledging the complexity of addiction can create space for healing that transcends both simplistic moral judgments and reductionist biological explanations. As Lieberman emphasizes in her analysis of biblical illness narratives, "the Bible functions as a catalogue of narratives about pain, illness, and suffering that, in the course of interpreting what is happening in the sufferer's life, communicate differing perceptions of illness and differing ethical visions of how the world should be." [24].

Strengths of the Classical Medical Model

The classical medical model offers several significant strengths in addressing addiction that physicians will readily recognize from their training and practice. Understanding these strengths allows clinicians to leverage them effectively while acknowledging their limitations.

The foundation in scientific rigor represents perhaps the most compelling advantage of the medical model for physicians. Grounded in empirical research, this approach provides objectively verifiable evidence regarding the neurobiological mechanisms of addiction and the efficacy of various treatments [7]. For physicians trained in evidence-based medicine, this scientific foundation offers a familiar framework for understanding addiction and evaluating treatment options. The ability to measure outcomes objectively, quantify treatment effects, and apply rigorous research methodologies aligns with fundamental principles of medical practice.

The medical model has directly contributed to medication development that has revolutionized addiction treatment. This framework has facilitated the development of pharmacotherapies that can significantly reduce withdrawal symptoms, block drug effects, diminish cravings, and support long-term recovery [32]. For physicians, medications like methadone, buprenorphine, naltrexone, and acamprosate represent concrete, prescribable interventions that can be integrated into treatment plans alongside psychosocial approaches. These pharmacological tools provide clinicians with tangible means to address the neurobiological aspects of addiction that often present significant barriers to recovery.

By framing addiction as a medical condition rather than a moral failing, the medical model has contributed substantially to destigmatization, a shift that physicians are uniquely positioned to advance through their clinical language and practice [33]. This reframing helps encourage treatment-seeking behavior among patients who might otherwise avoid care due to shame or fear of judgment. For physicians, adopting non-stigmatizing language and approaching addiction as they would any other medical condition creates a therapeutic alliance that supports effective treatment.

The medical model allows for addiction treatment to be integrated with mainstream healthcare systems, a significant advantage in an era of increasing recognition of addiction as a public health concern [34]. This integration potentially improves access, insurance coverage, and coordination with treatment for co-occurring medical and psychiatric conditions. For physicians practicing in integrated care settings, this systemic approach facilitates addressing the multiple health needs that often accompany substance use disorders.

Individualized treatment planning based on comprehensive assessment represents another strength of the medical model that aligns with contemporary medical practice [35]. The medical model emphasizes personalized treatment approaches tailored to the specific needs of each patient, including consideration of co-occurring disorders and genetic factors. This individualized approach parallels precision medicine efforts in other medical specialties and acknowledges the heterogeneity of substance use disorders.

Limitations of the Classical Medical Model

Despite its significant contributions, the medical model has notable limitations that physicians encounter in clinical practice when working with patients with substance use disorders. Recognizing these limitations allows for a more nuanced and comprehensive approach to treatment.

The tendency toward reductionism represents a significant limitation that physicians may observe in purely biomedical approaches to addiction. Critics argue that the medical model can reduce the complex phenomenon of addiction to neurobiological processes, potentially neglecting psychological, social, cultural, and spiritual dimensions that significantly influence both the development of addiction and recovery processes [36]. For physicians, this reductionist tendency may manifest as frustration when patients fail to respond to biologically-targeted interventions despite appropriate medication management. This limitation highlights the need for physicians to maintain awareness of the multidimensional nature of addiction even within a medical framework.

Over-medicalization presents another challenge that physicians must navigate carefully. Excessive focus on pharmacological solutions may underemphasize the importance of psychological

and social interventions, potentially leading to an over-reliance on medication that addresses symptoms without sufficient attention to underlying psychological processes or social determinants of health [37]. For physicians trained primarily in pharmacological interventions, this limitation underscores the importance of developing familiarity with complementary psychological and social approaches or establishing effective referral networks with mental health and social service providers.

The medical model's reliance on professional expertise, specialized facilities, and medications creates access barriers that physicians frequently encounter in practice [38]. These elements can be costly and inaccessible to many individuals, particularly those without adequate insurance or in underserved areas. For physicians working with marginalized populations or in resource-limited settings, these access barriers may significantly constrain treatment options and necessitate creative adaptations of the medical model to address local realities and available resources.

Compliance challenges represent a common limitation that physicians encounter when implementing medical treatments for addiction [39]. Medical treatments often face challenges with patient adherence, particularly when medications must be taken regularly over extended periods. For physicians, these challenges highlight the need for comprehensive patient education, motivational enhancement strategies, and attention to practical barriers to medication adherence such as cost, transportation, and side effects.

Perhaps most significantly for physicians interested in holistic care, the medical model may inadequately address existential aspects of addiction that significantly impact recovery outcomes [40]. Questions of meaning, purpose, and spiritual well-being that many individuals in recovery identify as crucial may receive insufficient attention in a purely biomedical approach. For physicians, this limitation suggests the importance of developing comfort with discussing existential concerns and recognizing when referral to spiritual care providers or mutual-help groups may complement medical treatment.

Strengths of the Recovery Model

The 12-Step approach offers distinctive advantages in addiction recovery that physicians can leverage to enhance treatment outcomes. Understanding these strengths allows clinicians to effectively incorporate 12-Step principles and participation into comprehensive treatment planning.

The accessibility and cost-effectiveness of 12-Step programs represent significant advantages that address common barriers to addiction treatment encountered in medical practice [41]. 12-Step meetings are widely available, free of charge, and accessible without professional referral or insurance, creating a low-barrier entry point for recovery support. For physicians concerned about treatment accessibility, particularly after discharge from acute care or when insurance coverage limits treatment duration, 12-Step participation offers a sustainable resource that extends beyond the constraints of the medical system. This accessibility is particularly valuable for patients in rural areas or those with limited financial resources.

The community and belonging fostered through 12-Step fellowships address social determinants of health that physicians increasingly recognize as critical to recovery outcomes [42]. The fellowship component provides social connection, reducing isolation and offering role models of successful recovery factors known to support sustained behavior change. For physicians, 12-Step communities represent a powerful adjunctive resource that addresses social factors difficult to target within traditional medical encounters. The peer support available through these communities offers validation, practical guidance, and accountability that complement professional interventions.

The spiritual dimension of the 12-Step approach addresses existential concerns that physicians may find challenging to address within typical medical encounters [43]. For many individuals, the spiritual framework addresses existential distress and provides a sense of meaning and purpose that supports recovery motivation. While physicians vary in their comfort discussing spiritual matters, recognizing the potential therapeutic value of spiritual engagement allows clinicians to support this dimension of recovery even when it falls outside their direct expertise. The spiritual framework offers a pathway to transcendence and meaning-making that many patients find essential to sustained recovery.

The 12-Step emphasis on comprehensive lifestyle change aligns with medical recognition of lifestyle factors in health outcomes [44]. The program encourages holistic transformation beyond abstinence, including improved relationships, ethical behavior,

and service to others. For physicians, this comprehensive approach addresses the broad lifestyle disruptions associated with addiction and supports the development of a healthy, non-substance-focused identity that protects against relapse. The focus on character development and relationship repair addresses psychological and social factors that often perpetuate substance use.

Unlike time-limited professional treatments constrained by insurance coverage or practice models, 12-Step programs offer long-term support throughout the recovery journey [20]. This ongoing support addresses the chronic nature of addiction and provides continuity that physicians cannot typically offer through medical visits alone. For physicians managing chronic conditions, the longitudinal support available through 12-Step participation offers reassurance that patients have access to recovery reinforcement between appointments and after formal treatment concludes.

Limitations of the Recovery Model

The 12-Step approach also presents several limitations that physicians should consider when integrating these programs into treatment plans. Understanding these limitations allows for more nuanced clinical recommendations and anticipation of potential barriers to patient engagement.

The spiritual emphasis central to the 12-Step approach can create significant barriers for some patients. The spiritual language and concept of a Higher Power may alienate individuals who are atheist, agnostic, or uncomfortable with spiritual frameworks [45]. For physicians working with diverse patient populations, this limitation necessitates careful consideration of each patient's worldview before strongly recommending 12-Step participation. Physicians should be prepared to discuss secular alternatives (such as SMART Recovery or Refuge Recovery) or to help patients interpret 12-Step concepts in ways compatible with their personal beliefs. This limitation also highlights the importance of avoiding assumptions about patients' receptivity to spiritually-oriented interventions.

The relatively standardized approach of 12-Step programs may inadequately address individual differences in addiction etiology, severity, co-occurring disorders, or cultural background [46]. While medical treatment increasingly moves toward precision medicine, the uniform program of 12-Step groups may not sufficiently accommodate the heterogeneity of substance use disorders. For physicians, this limitation underscores the importance of assessing fit between patient characteristics and recovery approaches rather than assuming universal applicability. Patients with significant psychiatric comorbidities, cognitive impairments, or cultural backgrounds that conflict with 12-Step norms may require alternative or additional supports.

Although research increasingly supports the effectiveness of 12-Step facilitation and participation for many individuals, physicians should note that the evidence base remains less extensive and rigorous than for some professionally delivered treatments [41]. This limitation reflects methodological challenges in researching mutual-help groups rather than ineffectiveness, but it may create hesitation among physicians trained in evidence-based medicine. For physicians navigating this limitation, familiarity with the growing body of research on 12-Step outcomes and mechanisms is beneficial, as is recognition that absence of evidence is not evidence of absence, particularly for approaches with low-risk profiles and potential benefits.

The emphasis on complete abstinence in most 12-Step programs may discourage harm reduction approaches that could benefit individuals not ready for or interested in abstinence-based recovery [47]. This absolute stance contrasts with medical approaches that increasingly recognize the value of harm reduction for patients unable or unwilling to achieve immediate abstinence. For physicians, this limitation suggests caution in recommending 12-Step programs to patients who explicitly reject abstinence goals or who might benefit from incremental harm reduction approaches as a pathway to eventual abstinence.

Certain concepts central to 12-Step programs, such as "powerlessness" and "character defects," can be misunderstood or misapplied in ways that may impede recovery for some individuals [48]. Without adequate context or skilled interpretation, these concepts might be experienced as demoralizing or stigmatizing by certain patients. For physicians, this limitation highlights the importance of discussing potential misinterpretations and providing clarification about 12-Step concepts when making referrals. Additionally, physicians should remain alert to signs that patients are experiencing 12-Step participation as detrimental rather than beneficial to their recovery process.



Rationale for Integration

The limitations of both models when applied in isolation suggest the potential value of an integrated approach that physicians can implement in clinical practice. Several compelling factors support this integration, creating opportunities for more comprehensive and effective addiction treatment.

The multifactorial nature of addiction represents perhaps the strongest rationale for integration. Substance use disorders involve complex interactions among biological, psychological, social, and spiritual dimensions that are best addressed through complementary approaches rather than single-model interventions [49,50]. For physicians, this complexity necessitates treatment strategies that address neurobiological mechanisms while

simultaneously attending to psychological processes, social contexts, and existential concerns. Neither the medical model nor the 12-Step approach alone fully encompasses this complexity, but together they offer complementary perspectives that more completely address the multidimensional nature of addiction.

A growing body of research provides evidence for the effectiveness of combined approaches that integrate medical and mutual-help elements. Studies indicate that outcomes improve when pharmacological and psychosocial interventions are combined, and when professional treatment is followed by mutual-help participation [51,52]. For physicians committed to evidence-based practice, this research supports integration as a means of optimizing patient outcomes rather than as merely a theoretical preference. The sequential integration of medical interventions followed by ongoing 12-Step participation has demonstrated particular effectiveness in maintaining treatment gains over time.
Contemporary healthcare systems increasingly recognize the need for recovery-oriented systems of care that extend beyond acute treatment to support long-term recovery [53,54]. These systems acknowledge addiction as a chronic condition requiring ongoing management rather than an acute problem resolved through time-limited interventions. For physicians practicing within these evolving systems, integration of medical and 12-Step approaches aligns with the shift toward longitudinal, person-centered care that addresses both immediate symptoms and long-term recovery maintenance. This integration bridges the gap between acute medical interventions and ongoing recovery support.

The reality of diverse recovery pathways further supports integration as a clinical approach. Research increasingly demonstrates that individuals recover through various pathways, suggesting the value of flexible approaches that can be tailored to personal needs, preferences, and cultural contexts [55]. For physicians, this diversity highlights the importance of offering multiple pathways to recovery rather than adhering rigidly to single-model approaches. Integration allows clinicians to honor patient autonomy by presenting complementary options that patients can engage with based on their individual circumstances and preferences.

Principles of Integration

Effective integration of the medical and 12-Step models requires adherence to several key principles that can guide physician practice. These principles offer a framework for thoughtfully combining approaches rather than simply layering them without conceptual coherence. Complementarity represents a fundamental principle for integration, recognizing that the models address different aspects of addiction and recovery. The medical model primarily targets biological mechanisms of addiction, while the 12-Step approach addresses spiritual and community dimensions [56]. For physicians, this complementarity suggests approaching integration not as an either/or choice but as a both/and opportunity. Medical interventions can address neurobiological aspects of addiction while 12-Step participation simultaneously addresses experiential and communal dimensions. This complementary approach acknowledges that different components of addiction require different intervention strategies.

Integration can occur through both sequential and concurrent application of medical interventions and 12-Step principles. Sequential integration might involve medical detoxification followed by 12-Step facilitation, while concurrent integration could combine medication management with simultaneous 12-Step participation [52]. For physicians, this principle offers flexibility in timing and combining interventions based on patient needs and treatment phase. Acute withdrawal might necessitate primarily medical approaches, while post-acute care might emphasize 12-Step engagement alongside continued medication management. This principle recognizes that the optimal balance shifts throughout the recovery process.

Personalization remains essential in integrated approaches, as the optimal balance of medical and 12-Step elements will vary significantly among individuals [55]. While integration offers a more comprehensive approach than either model alone, the specific integration must be tailored to individual needs, preferences, and circumstances. For physicians, this principle translates to individualized treatment planning that accounts for factors such as addiction severity, co-occurring conditions, previous recovery experiences, spiritual orientation, and cultural background. This personalization honors patient autonomy while optimizing outcomes through targeted intervention.

Cross-model communication between medical providers and 12-Step communities fosters effective integration by overcoming historical tensions and creating coordinated recovery support [57]. For physicians, this principle encourages developing familiarity with local 12-Step resources, understanding 12-Step terminology and concepts, and maintaining open dialogue with patients about their 12-Step experiences. This communication bridges potential gaps between professional treatment and mutual-help participation, creating continuity rather than fragmentation in the recovery process.

Throughout integration efforts, respect for patient autonomy in choosing recovery pathways remains paramount [58]. Integration should avoid coercion into either model or should honor patient preferences even when these diverge from provider recommendations. For physicians, this principle translates to presenting options, providing information about potential benefits and limitations of different approaches, and supporting patient decision-making. This respect acknowledges both the physician's expertise and the patient's fundamental right to determine their recovery path based on personal values and circumstances.

Practical Applications

The integrated framework can be implemented through various practical approaches that physicians can incorporate into their clinical practice. These applications translate theoretical integration into tangible interventions that improve patient care.

Physicians can provide medically-supported 12-Step participation by prescribing medication for withdrawal management, craving reduction, or psychiatric comorbidities while actively supporting patient engagement in 12-Step programs. This support extends beyond mere referral to include education about 12-Step principles, encouragement of attendance, and facilitated connections to local meetings [59]. In practice, this might involve physicians discussing how medications like naltrexone or acamprosate complement rather than conflict with 12-Step participation, addressing potential stigma patients might encounter regarding medication use in 12-Step contexts, and scheduling follow-up discussions about 12-Step experiences during medication management appointments.

Healthcare providers can incorporate elements of 12-Step facilitation therapy into clinical practice, helping patients understand and engage with 12-Step principles in ways compatible with their medical treatment [60]. This facilitation might include explaining 12-Step concepts using language aligned with medical understandings, helping patients identify meetings appropriate to their needs and preferences, and addressing barriers to participation. For physicians, this application might involve brief motivational interventions during routine appointments that encourage 12-Step attendance or preparing patients for what to expect at their first meeting. These facilitation elements can be integrated into standard clinical encounters without requiring extensive additional time.

Long-term monitoring and support through recovery management checkups can combine medical assessment with encouragement of mutual-help participation, addressing the chronic nature of addiction [61]. These regular check-ins acknowledge that recovery extends beyond acute treatment episodes and requires ongoing attention similar to other chronic conditions. For physicians, this approach might involve scheduling periodic follow-up appointments specifically focused on recovery maintenance rather than only crisis intervention. During these appointments, physicians can monitor both medical aspects of recovery and 12-Step participation, adjusting the integrated treatment plan as needed based on recovery progress or challenges.

Formally integrating peer recovery specialists often individuals in personal recovery through 12-Step programs into medical treatment settings can bridge the gap between professional care and mutual-help communities [62]. These specialists provide experiential knowledge that complements clinical expertise and can facilitate connections to community resources. For physicians, collaboration with peer specialists might involve incorporating them into treatment teams, consulting them regarding patient barriers to 12-Step engagement, or co-leading patient education sessions that combine medical and experiential perspectives on recovery. This integration brings authentic recovery experience into medical settings in a structured manner.

Modified versions of the 12 Steps that maintain core principles while using language more compatible with medical or secular perspectives can increase accessibility for patients who might otherwise resist 12-Step participation [20]. These adaptations preserve essential recovery mechanisms while reducing barriers related to terminology or conceptual frameworks. For physicians, awareness of these adaptations (such as SMART Recovery or secular 12-Step interpretations) provides additional options for patients who express discomfort with traditional 12-Step language or concepts. Physicians can discuss these alternatives with patients who might benefit from mutual-help participation but resist conventional 12-Step approaches.



Case History

Integrated Treatment for Alcohol Use Disorder

To illustrate the integrated framework, consider the following case example that demonstrates how physicians might implement integration in clinical practice:

John, a 45-year-old executive, presented to his primary care physician with elevated liver enzymes, hypertension, and symptoms of depression. After screening and assessment reveal severe alcohol use disorder, his physician initiated an integrated treatment plan that addressed both medical and recovery dimensions:

He was prescribed the usual cocktail of medication for alcohol withdrawal symptoms and was referred to an addiction specialist who prescribed naltrexone to reduce cravings and prevent relapse. This pharmacological approach directly addressed the neurobiological aspects of John's alcohol use disorder, reducing withdrawal discomfort and diminishing the reinforcing effects of alcohol. He was prescribed these medications as tools that created a physiological foundation for recovery work rather than as complete solutions in themselves.

Recognizing the frequent co-occurrence of mood disorders and alcohol use disorders, the physician arranged for psychiatric evaluation. A psychiatrist evaluated John's depression and prescribed an antidepressant compatible with his addiction treatment, carefully selecting a medication with low abuse potential and beneficial interactions with naltrexone. This attention to psychiatric comorbidity acknowledges the complex interplay between mood and substance use that requires specific medical attention within the integrated framework.

John underwent cognitive-behavioral therapy to address maladaptive thought patterns and to develop coping skills for managing stress and cravings. This evidence-based psychological intervention complements the biological focus of medication by

addressing psychological mechanisms that maintain addictive behavior. The therapist coordinates with John's physicians to ensure alignment between psychological and pharmacological approaches, creating coherence rather than fragmentation in his treatment experience.

In parallel with these medical interventions, John's therapist provided 12-Step facilitation, explaining the principles of AA and encouraging attendance. Rather than presenting 12-Step participation as an alternative to medical treatment, the therapist framed it as a complementary approach that addresses dimensions of recovery not fully covered by medical interventions. The facilitation includes practical guidance about meeting options, discussion of potential barriers to participation, and clarification of commonly misunderstood concepts. With this preparation, John began attending AA meetings, obtains a sponsor, and starts working through the 12 Steps. His engagement with the fellowship provided peer support, accountability, and a framework for personal growth that extended beyond his formal treatment sessions. The mutual-help community offered continuous support between professional appointments and introduced John to role models who demonstrate long-term recovery, providing hope and practical guidance from lived experience.

The integration continued through ongoing coordination among treatment providers. Regular appointments with his healthcare providers include discussions of his 12-Step experiences, with clinicians expressing interest in his fellowship participation and helping him navigate any challenges encountered. His sponsor, educated about the medical aspects of addiction, was supportive of his medication use, recognizing it as part of his recovery plan rather than as a substitute for spiritual growth or fellowship participation.

After completing the acute phase of treatment, John participated in recovery monitoring through regular medical check-ups that assess both physiological markers of health and recovery status. He continues AA attendance after formal treatment ends, illustrating the integration of medical care and mutual-help support in long-term recovery management. His physician maintains naltrexone while gradually extending the interval between appointments based on recovery stability, demonstrating the dynamic nature of integrated care over time.

This case demonstrates how medical and 12-Step approaches can be synergistically combined to address the biological, psychological, social, and spiritual dimensions of addiction. The integration provides comprehensive support that neither approach alone could offer, creating a recovery framework that adapts to changing needs throughout the recovery process.

Conclusion

The classical medical model and the 12-Step recovery model represent distinct but potentially complementary approaches to understanding and addressing addiction. For physicians navigating the complex landscape of addiction treatment, the integration of these models offers a promising framework that addresses limitations inherent in either approach applied in isolation.

The medical model's considerable strengths in addressing biological aspects of addiction through evidence-based interventions provide a foundation for recovery that many patients require. Neurobiological insights regarding reward circuitry, withdrawal mechanisms, and medication effects offer physicians powerful tools to address the physiological dimensions of addiction that can present significant barriers to recovery initiation. Yet as many clinicians observe in practice, these biological interventions alone often prove insufficient for sustainable recovery, particularly given the complex psychosocial and spiritual dimensions of addiction that extend beyond neurochemistry. The medical model, while necessary, is rarely sufficient for comprehensive addiction care.

The 12-Step model offers complementary strengths through its emphasis on spiritual transformation, community support, and ongoing recovery maintenance dimensions that medical interventions alone typically cannot address. The fellowship component provides social connection and accountability that extend beyond the temporal and practical constraints of medical appointments, while the spiritual framework addresses existential questions that many patients identify as central to their recovery process. These elements address aspects of addiction that resist purely medical intervention yet significantly impact treatment outcomes and quality of life in recovery.

An integrated framework acknowledges addiction as a complex condition requiring multifaceted approaches rather than single-model interventions. By thoughtfully combining elements of both models based on individual needs and preferences, physicians can offer more comprehensive care that addresses the full spectrum of factors involved in addiction and recovery. This integration respects both the biological reality of addiction and the lived experience of recovery that encompasses psychological, social, and spiritual dimensions. It allows physicians to maintain scientific rigor while embracing complementary approaches that enhance treatment effectiveness.

The practical applications of integration discussed in this paper medically-supported 12-Step participation, 12-Step facilitation in medical settings, recovery management checkups, peer recovery support integration, and adapted 12-Step approaches offer physicians tangible strategies for implementation. These applications translate theoretical integration into clinical practice that can be adapted to various healthcare settings and patient populations. The case example illustrates how these strategies can be combined to create coherent, patient-centered care that addresses multiple dimensions of addiction simultaneously.

For physicians committed to evidence-based practice, continued research on integration methods remains essential. Future research should explore optimal methods for integration, including identifying which individuals benefit most from various combinations of medical and 12-Step approaches, developing structured protocols for integration, and examining outcomes of integrated care in diverse populations and settings. This research

will further refine integration strategies and strengthen the evidence base for comprehensive addiction treatment.

In conclusion, the integration of medical and 12-Step approaches offers physicians a more complete framework for addressing the complex phenomenon of addiction. By recognizing the complementary strengths of each model and implementing thoughtful integration strategies, clinicians can provide care that honors both scientific understanding and human experience. This integration represents not a compromise between approaches but rather a synergy that enhances the effectiveness of addiction treatment and supports sustainable recovery for patients with substance use disorders.

References

1. Porter R. The Cambridge history of medicine. Cambridge: Cambridge University Press. 2006.
2. Bynum WF. History of medicine: A very short introduction. Oxford: Oxford University Press. 2008.
3. White WL. Slaying the dragon: The history of addiction treatment and recovery in America. Bloomington, IL: Chestnut Health Systems. 2014.
4. Katcher BS. Benjamin Rush's educational campaign against hard drinking. Am J Public Health. 1993; 83 :273-81.
5. White WL. Slaying the dragon: The history of addiction treatment and recovery in America. Bloomington, IL: Chestnut Health Systems. 1998.
6. Leshner AI. Addiction is a brain disease, and it matters. Science. 1997; 278: 45-7.
7. Volkow ND, Koob GF, McLellan AT, et al. Neurobiologic advances from the brain disease model of addiction. N Engl J Med. 2016; 374: 363-71.
8. Kurtz E. Not-God: A history of Alcoholics Anonymous. Center City, MN: Hazelden. 1979.
9. White WL, Kurtz E, Galanter M, et al. Twelve defining moments in the history of Alcoholics Anonymous. Recent dev alcohol. 2008; 18: 37-57.
10. Alcoholics Anonymous. Alcoholics Anonymous: The story of how many thousands of men and women have recovered from alcoholism. 4th ed. New York: Alcoholics Anonymous World Services. 2001.
11. Kurtz E. Alcoholics Anonymous and the disease concept of alcoholism. Alcohol Treat Q. 2002; 20: 5-39.
12. Anderson DJ, McGovern JP, DuPont RL, et al. The origins of the Minnesota model of addiction treatment a first person account. J Addict Dis. 1999; 18: 107-14.
13. McElrath D. The Minnesota model. J Psychoactive Drugs. 1997; 29: 141-4.
14. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th ed. Arlington, VA: American Psychiatric Publishing. 2013.
15. Carroll KM, Kiluk BD. Cognitive behavioral interventions for alcohol and drug use disorders: Through the stage model and back again. Psychol Addict Behav. 2017; 31: 847-61.
16. Humphreys K, McLellan AT. A policy-oriented review of strategies for improving the outcomes of services for substance use disorder patients. Addiction. 2011; 106: 2058-66.
17. McLellan AT, Lewis DC, O'Brien CP, et al. Drug dependence, a chronic medical illness: Implications for treatment, insurance, and outcomes evaluation. JAMA. 2000; 284: 1689-95.
18. Tonigan JS, Rynes KN, McCrady BS, et al. Spirituality as a change mechanism in 12-step programs: A replication, extension, and refinement. Subst Use Misuse. 2013; 48: 1161-73.
19. Kelly JF. Is Alcoholics Anonymous religious, spiritual, neither? Findings from 25 years of mechanisms of behavior change research. Addiction. 2017; 112: 929-36.
20. Humphreys K. Circles of recovery: Self-help organizations for addictions. Cambridge: Cambridge University Press. 2004.
21. Kelly JF, Magill M, Stout RL, et al. How do people recover from alcohol dependence? A systematic review of the research on mechanisms of behavior changes in Alcoholics Anonymous. Addict Res Theory. 2011; 17: 236-59.
22. Lieberman R. Sin or sickness? The role of divine punishment in conceptualizing addiction. J Spiritual Health. 2019; 27: 412-28.
23. Lieberman R. Divine retribution and medical explanation: Historical tensions in addiction treatment frameworks. J Med Humanities. 2020; 41: 178-92.
24. Liebermann R. Is illness a divine punishment? "I YHWH am your healer" Exodus 15:26. TABS: The Torah and Biblical Scholarship. 2023. https://thetorah.com/article/is-illness-a-divine-punishment.
25. Avrahami Y. The senses of scripture: Sensory perception in the Hebrew Bible. New York: T&T Clark. 2012.
26. Ungar-Sargon J. The neurobiology of existential distress: Implications for addiction medicine. J Addict Neurosci. 2019; 12: 214-29.
27. Ungar-Sargon J, Mikhailova T. Existential anxiety and substance cravings: A dual-process neuroimaging study. Neuropsychopharmacology. 2020; 45: 1392-1401.
28. Ungar-Sargon J. Bridging the divide: Toward a neurobiologically-informed approach to existential medicine. Adv Mind Body Med. 2021; 35: 8-17.
29. Lieberman R. Language of punishment in addiction treatment discourse: A critical discourse analysis. Health Commun. 2021; 36: 1104-16.
30. Ungar-Sargon J, Damasio A, Phillips AG, et al. Parasympathetic responses to surrender: Autonomic correlates of 12-step participation. Front Psychiatry. 2022; 13: 784926.
31. Ungar-Sargon J. Neurobiological-existential integration in addiction treatment: Clinical applications for physicians. JAMA Psychiatry. 2023; 80: 557-568.
32. Kampman K, Jarvis M. American Society of Addiction Medicine (ASAM) National Practice Guideline for the use of medications in the treatment of addiction involving opioid use. J Addict Med. 2015; 9: 358-67.

33. Kelly JF, Westerhoff CM. Does it matter how we refer to individuals with substance-related conditions? A randomized study of two commonly used terms. Int J Drug Policy. 2010; 21: 202-7.
34. McLellan AT, Woodworth AM. The affordable care act and treatment for "substance use disorders": Implications of ending segregated behavioral healthcare. J Subst Abuse Treat. 2014; 46: 541-5.
35. Woody GE, Blaine J, DuPont RL, et al. Depression in narcotic addicts: Quite possibly more than a chance association. Handbook on drug abuse. Washington, DC: National Institute on Drug Abuse. 1979; 277-85.
36. Heather N. Q: Is addiction a brain disease or a moral failing? A: Neither. Neuroethics. 2017; 10: 115-24.
37. Hall W, Carter A, Forlini C, et al. The brain disease model of addiction: Is it supported by the evidence and has it delivered on its promises? Lancet Psychiatry. 2015; 2: 105-10.
38. Wakeman SE, Larochelle MR, Ameli O, et al. Comparative effectiveness of different treatment pathways for opioid use disorder. JAMA Netw Open. 2020; 3: e1920622.
39. Swift RM, Kallio M, Rohsenow DJ, et al. Medication treatment for alcohol use disorders. Addiction recovery management: Theory, research and practice. New York: Humana Press; 2011: 287-309.
40. Galanter M. Spirituality and recovery in 12-step programs: An empirical model. J Subst Abuse Treat. 2007; 33: 265-72.
41. Kelly JF, Humphreys K, Ferri M, et al. Alcoholics Anonymous and other 12-step programs for alcohol use disorder. Cochrane Database Syst Rev. 2020; 3: CD012880.
42. Moos RH. Active ingredients of substance use-focused self-help groups. Addiction. 2008; 103: 387-96.
43. Galanter M, Dermatis H, Post S, et al. Spirituality-based recovery from drug addiction in the Twelve-Step fellowship of Narcotics Anonymous. J Addict Med. 2013; 7: 189-95.
44. Kelly JF, Greene MC. Beyond motivation: Initial validation of the Commitment to Sobriety Scale. J Subst Abuse Treat. 2014; 46: 257-63.
45. Tonigan JS, Miller WR, Schermer C, et al. Atheists, agnostics and Alcoholics Anonymous. J Stud Alcohol. 2002; 63: 534-41.
46. Ferri M, Amato L, Davoli M, et al. Alcoholics Anonymous and other 12-step programmes for alcohol dependence. Cochrane Database Syst Rev. 2006; 3:CD005032.
47. Marlatt GA, Witkiewitz K. Update on harm-reduction policy and intervention research. Annu Rev Clin Psychol. 2010; 6: 591-606.
48. Kelly JF, Hoeppner B, Stout RL, et al. Determining the relative importance of the mechanisms of behavior change within Alcoholics Anonymous: A multiple mediator analysis. Addiction. 2012; 107: 289-99.
49. Engel GL. The need for a new medical model: A challenge for biomedicine. Science. 1977; 196: 129-36.
50. Kaskutas LA, Borkman TJ, Laudet A, et al. Elements that define recovery: The experiential perspective. J Stud Alcohol Drugs. 2015; 76: 954-63.
51. Weiss RD, Griffin ML, Gallop RJ, et al. The effect of 12-step self-help group attendance and participation on drug use outcomes among cocaine-dependent patients. Drug Alcohol Depend. 2000; 60: 271-80.
52. Humphreys K, Blodgett JC, Wagner TH, et al. Estimating the efficacy of Alcoholics Anonymous without self-selection bias: An instrumental variables re-analysis of randomized clinical trials. Alcohol Clin Exp Res. 2014; 38: 2688-94.
53. White WL. Recovery management and recovery-oriented systems of care: Scientific rationale and promising practices. Pittsburgh, PA: Northeast Addiction Technology Transfer Center. 2008.
54. Kelly JF, White WL. Addiction recovery management: Theory, research and practice. New York: Humana Press. 2011.
55. Kelly JF, Greene MC, Bergman BG, et al. Beyond abstinence: Changes in indices of quality of life with time in recovery in a nationally representative sample of U.S. adults. Alcohol Clin Exp Res. 2018; 42: 770-80.
56. Miller WR, Kurtz E. Models of alcoholism used in treatment: Contrasting AA and other perspectives with which it is often confused. J Stud Alcohol. 1994; 55: 159-66.
57. White WL, Boyle M, Loveland D, et al. Alcoholism/addiction as a chronic disease: From rhetoric to clinical reality. Alcohol Treat Q. 2006; 20: 107-29.
58. Miller PG, Miller WR. What should we be aiming for in the treatment of addiction? Addiction. 2009; 104: 685-6.
59. Timko C, DeBenedetti A, Billow R, et al. Intensive referral to 12-Step self-help groups and 6-month substance use disorder outcomes. Addiction. 2006; 101: 678-88.
60. Project MATCH Research Group. Matching alcoholism treatments to client heterogeneity: Project MATCH posttreatment drinking outcomes. J Stud Alcohol. 1997; 58: 7-29.
61. Dennis ML, Scott CK. Managing addiction as a chronic condition. Addict Sci Clin Pract. 2007; 4: 45-55.
62. Bassuk EL, Hanson J, Greene RN, et al. A. Peer-delivered recovery support services for addictions in the United States: A systematic review. J Subst Abuse Treat. 2016; 63: 1-9.

© 2025 Ungar-Sargon J. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License