

## Evaluating the Therapeutic Effect of Vitamin C on the Biological Damages of the Low Electromagnetic Fields in Wistar Mice

Ali Neamati<sup>1</sup>, Ghorban Safaeian layen<sup>2\*</sup> and Zahra Safaeian layen<sup>3</sup>

<sup>1</sup>Department of Biochemistry and Biophysics, Mashhad Branch, Islamic Azad University, Iran.

<sup>2</sup>Department of Technology of Radiology, School of Paramedical science, Mashhad University of Medical Science, Mashhad, Iran

<sup>3</sup>Department of Medical Laboratory Sciences, School of Paramedical Science, Mashhad University of Medical Science, Mashhad, Iran.

### \*Correspondence:

Ghorban Safaeian Layen, Department of Technology of Radiology, School of Paramedical Science, Mashhad University of Medical Sciences, Mashhad, Iran.

Received: 13 Jun 2022; Accepted: 20 Jul 2022; Published: 25 Jul 2022

**Citation:** Neamati A, layen GS, layen ZS. Evaluating the Therapeutic Effect of Vitamin C on the Biological Damages of the Low Electromagnetic Fields in Wistar Mice. J Med - Clin Res & Rev. 2022; 6(7): 1-6.

### ABSTRACT

*With the development of advanced telecommunication systems such as cell phones which are used by more than 75% of the world population, nonionizing electromagnetic waves are believed to be among the probable reasons for the increasing number of cases diagnosed with different types of cancers, depression, and hormonal disorders and the growing rate of unwanted abortions.*

*This study sets out to examine whether taking Vitamin C and doing short-term aerobic exercises could be effective in reducing (up to 65-75% VO<sub>2</sub>MAX) such harmful effects. In this experimental study, 24 adult male mice (Balb/C) were divided into 3 groups. The control group consist of 8 mice were kept in normal conditions. The group exposure to Electromagnetic field consists of 8 mice, which were exposed to ELF (50Hz, 4Mt) for 15 days (4 hours per day), and the group that exposure to Electromagnetic field supplemented with vitamin C, contains 8 mice that were exposed to ELF (50Hz, 4Mt) for 15 days (4 hours per day) and 0.5 cc vitamin C were injected intraperitoneally 7 times. This injection was done 3 times a week. After 15 days, mice were anesthetized by chloroform, following careful dissection, organs were splitted over then this samples were dissected and measured. Data was analyzed using one-way ANOVA test.*

*The results of the unilateral variance analysis administered on the collected data show a significant difference among the groups set in the study, indicating that 15 days of exposure to electromagnetic waves (mobile frequency) caused some disorders. The Results also indicate significant differences between pre-test and post-test of all the variables defined for the study.*

### Keywords

Electromagnetic waves, Radiation, Aerobic exercises, VO<sub>2</sub>MAX, body weight, Vitamin C.

### Introduction

During last years, increased use of devices producing electromagnetic wave has attracted researcher's attention to possible effects of them on human health. Electromagnetic waves have different shapes, they are classified according to their frequencies and wave length [1]. Today, effects of electromagnetic

waves with low frequency (ELF) have been attracted researcher's attention [2,3]. Electromagnetic waves with different intensities influence on prevalence of embryo evolution disorders, infertility, neuronal/sleep disorders, gastro-intestinal, heart disorders and several neoplasms including hematopoietic, lymphoma [4,5].

Thymus is behind of sternum between lungs, often it contains 2 parts. Each part of thymus is divided to smaller parts. Cortex contains compacted lymphocytes. Medulla constructed from sporadic epithelial and lymphocytes. Epithelial cells in thymus

produce its hormones; possibly, it incorporates in T lymphocyte evolution; however, its effect is not clear. Thymus is large in childhood but during growth, it will be replaced by connective and adipose tissues. During maturation, thymus will be smaller but it incorporates in T cell evolution. Thymus produces many hormones like thymosin, thymic humoral factor, thymopoietin; mentioned hormones induce proliferation and differentiation of T cells. Some evidence reveals that thymus hormones can delay aging [6]. Testis produces hormones and spermatozoon. Testes are wrapped by tunica albuginea. Tunica albuginea thickens at posterior of testis and constructs mediastinum, which is a fibrous connective tissue, diffuses to gland and divides it to pyramid structures called testis lobules. These walls are not complete; often there are connections between them. Each lobule contains 1-4 seminiferous tubules. They locate in a fragile connective membrane, which is full of blood vessels, lymphocytes, neurons and leydig cells. Seminiferous tubules produce male reproductive cells, spermatozoon. While leydig cells produce androgenic hormones, sperms are produced in seminiferous tubules. Seminiferous tubules are connective tissues contain a basal and a germ epithelial layer. Fibrous tunica propria encompasses seminiferous tubules and has several fibroblast layers. Innermost layer attached to basal layer is constructed from semi-muscular flat cells. These cells function as smooth muscles. Leydig cells occupy most part of space between seminiferous tubules. Epithelial seminiferous tissue has 2 cells; sertoli or supportive cells and cells which construct spermatogene antecedent cells. Spermatogene antecedent cells are distributed in 4-8 layers, they produce spermatozoids. Process of producing spermatozoids is called spermatogenesis. This includes cells divisions via mitosis/meiosis and final differentiation of spermatozoids, which is called spermiogenesis [7].

Vitamin C or L-ascorbate is a vital micronutrient for many species like human, monkeys and little number of other mammals, especially guinea pigs and some bird/fish species. Amount of vitamin C is a critical index for commercial value of fruits and vegetables. Vitamin C is a solid, white, water soluble, safe component, which has a circular ester; it is hydrolyzed in aqueous medium and convert to acid [8]. This vitamin, which attacks to electrons in body reactions, is most important antioxidant in neutralizing free radicals and inhibiting oxidative stress [9].

Antioxidant mechanism of action: free radical/reactive oxygen species production is an inevitable issue in metabolism process. These compounds because of high potential in damaging biological macromolecules like fats, DNAs function as major agents of ageing and cause several diseases. Organisms have various mechanisms to defend and neutralize these reactive species including defensive anti-oxidant immune systems. These systems have enzymes like superoxide dismutase, catalase. Glutathione peroxidase and macro molecules like albumin, ceruloplasmin, ferritin and micro molecules like carotene, alpha-tocopherol, ascorbate, methionine, uric acid, bilirubin, and reduced glutathione (GSH) [10].

There are several reports about role of this vitamin in prevention DNA damages and occurrence cancer or heart diseases [11]. Large

amount of vitamin C aggregates in ovaries and other endocrine tissues. In ovaries, vitamin C aggregation occurs in granulosa, luteal and theca cells, these cells relate closely to fertility. Studies on luteal granulosa cells demonstrated that vitamin C can increase progesterone production/concentration [12]. This relation is strong negative feedback, where high levels of progesterone reduce vitamin C concentration and its metabolism is inhibited [9]. Fritze et al. exposed rats to 900 Hz electromagnetic waves, they observed that these waves increased Hsp7z mRNA replication at brain cortex [13]. In Louis studies, pregnant CD-1 mice exposed to 2.45 GHz waves for 100 minutes from 1-17 pregnancy days, rats were dissected at 18<sup>th</sup> day and their embryos were evaluated in terms of disorders. Results showed that weight of experimental embryos was less than control [14]. In another study, few Spagu-Dawley rats were exposed to microwaves from 6-20 pregnancy day. Weight of test group was less than control [15]. Oconnor exposed mice to 2450 MHz and found that weight of embryos was reduced significantly because of maternal thermal stress [16]. Kolomyitseva et al. investigated muridae, they reported that unionizing electromagnetic waves caused lipid storage in adipose tissue and weight gain [17]. Results of several studies on different effects of electromagnetic waves reveal that change of field intensity, even milli tesla has various biological effects; therefore, investigation on electromagnetic effects can be done in very wide range [18]. In this research, amount of these effects on biomass, thymus, testis and therapeutic role of vitamin C in improvement of changes caused by electromagnetic field, have been studied.

## Material and Methods

This experiment was done in research electrophysiology laboratory of biochemistry-biophysics department of Mashhad Azad University. In order to evaluate effects of 50 Hz electromagnetic waves on biomass, testis, thymus and therapeutic effect of vitamin C on mature male mice, 3-2.5 month Balb/C mice with average weight 20-25 grams were used. Mature mice have been stored in animal room with 60-70% humidity,  $23 \pm 1$  centigrade temperature and 12 hours light/12 hours overnight. Automatic electrical timer adjusted room light alternatively and radiators and cooler controlled temperature at winter and summer respectively. Cages were cleansed every other day. For smoothness of cage, wood chips were used. Mice feed was special prepared pellets (Khorasan Javaneh dam company), water delivered by flasks.

Mice were purchased from Razi Vaccine and Serum research Institute of Mashhad, for being adapted with new environment and eliminating stress of changing environment, mice were kept one week in animal room before experiment. Electromagnetic producing machine includes 35 cm diameter and 60 cm length PVC tube equipped with 1900 copper coils, which coiled 3 times around tube. This machine could produce 0.5-4 mT/25-100 Hz electromagnetic field. In order to evaluate therapeutic role of vitamin C, commercial 250 Osveh vitamin C tablets (chewable-scored tablets) were used. In order to inject tablet, it was dissolved in 50 cc physiologic serum. Intra-proteanal vitamin C was injected by disposable syringe. Study groups in this 15 day research were


15 day old ( 24 mature 2.5-3 month mice) distributed in 3 groups: 1) control: 8 mice stored in animal room with normal conditions, 2) exposed to electromagnetic waves group: in this group, 8 mice exposed to 50 Hz/4 mT electromagnetic waves for 15 days, 4 hours per day(12-16 hours), 3) exposed to electromagnetic waves receiving vitamin C: in this group, 8 mice exposed to 50 Hz/4 mT electromagnetic waves for 15 days whom were treated with 0.5 cc intra peritoneal vitamin C 7 times. This injection was done weekly, 3 times per week. During experiment, mice were weighted 3 times weekly by 0.01 resolution scale and their weights noted. In order to be sure about weighting, this process repeated 3 times. After required time, mice were anesthetized by chloroform, following careful dissection, organs split over. Then these organs exposed to physiologic serum for 3 seconds, dried on paper for 3 seconds. Organs were located separately on glass to weight. Spss16 did statistical analysis and Excel illustrated plots. Data were represented by mean  $\pm$  SEM. Comparison between control and test was done by statistical T-test and  $P < 0.05$  was considered statistically significant difference.




**Figure 1:** Weighing of Mice.



**Figure 2:** Dissection of Mice.



**Figure 3:** Weighing Mice Organs.




**Figure 4:** Intra-Protaneal Vitamin C Injection.

## Results

### Evaluation of Thymus Weight

Mean weight of thymus in control group is  $0.055 \pm 0.0120$  and in exposed electromagnetic fields is  $0.115 \pm 0.0066$ , there is significant difference  $P < 0.01$  between them. Mean weight of thymus in exposed electromagnetic receiving vitamin C group

is  $0.0131 \pm 0.0081$  which is not significant difference to exposed field group.




**Diagram 1:** Comparison Weight of Thymus Between Control, Electromagnetic Field Exposed and Electromagnetic Field Exposed Receiving Vitamin C Groups.

++ refers to significant difference between control and electromagnetic field exposed groups with  $P < 0.01$ .

NS refers to un-significant statistical difference.

#### Evaluation of Testis Weight

Mean weight of testis in control group is  $0.0916 \pm 0.0079$  and in exposed electromagnetic fields is  $0.045 \pm 0.0104$ , there is significant difference  $P < 0.01$  between them. Mean weight of testis in exposed electromagnetic receiving vitamin C group is  $0.068 \pm 0.0102$  which is not significant difference to exposed field group.




**Diagram 2:** Comparison Weight of Testis Between Control, Electromagnetic Field Exposed and Electromagnetic Field Exposed Receiving Vitamin C Groups.

++ refers to significant difference between control and electromagnetic field exposed groups with  $P < 0.01$ .

NS refers to un-significant statistical difference.

#### Evaluation of Body Weight

Mean weight of body in control group is  $33.615 \pm 0.8700$  and in exposed electromagnetic fields is  $23.542 \pm 0.4068$ , there is significant difference  $P < 0.001$  between them. Mean weight of testis in exposed electromagnetic receiving vitamin C group is  $26.422 \pm 0.8551$  which is significant difference to exposed field group with  $P < 0.05$ .



**Diagram 3:** Comparison Weight of Body Between Control, Electromagnetic Field Exposed and Electromagnetic Field Exposed Receiving Vitamin C Groups.

++ refers to significant difference between control and electromagnetic field exposed groups with  $P < 0.001$ .

\* Refers to significant statistical difference between electromagnetic field exposed and electromagnetic field Balb/c male mice exposed receiving vitamin C groups with  $P < 0.05$ .

#### Discussion and Conclusion

**Analysis the Results of 50 Hz/4 Mt Electromagnetic Wave Effects on Thymus Weight of Mature Balb/C Male Mice:** According to this research, 50 Hz/4 mT electromagnetic waves reduce thymus weight of mature Balb/C male mice which shows significant relation.

**Analysis the Results of 50 Hz/4 Mt Electromagnetic Effects on Thymus Weight of Mature Male Balb/C Mice Receiving Vitamin C:** According to this research, 50 Hz/4 mT electromagnetic waves did not change significantly thymus weight of mature male Balb/C mice receiving vitamin C.

**Analysis the Results of 50 Hz/4 Mt Electromagnetic Wave Effects on Testis Weight of Mature Balb/C Male Mice:** According to this research, 50 Hz/4 mT electromagnetic waves reduce testis weight of mature Balb/C male mice which shows significant relation.

Studies on total weight of testis and epididymis showed significant weight reduction after exposure to electromagnetic field, which is in agreement with current, study and reveals the discharge of sperms at seminiferous tubules or epididymis; because a semi-muscular layer surrounds seminiferous layer in mice, which has erectile property. Electromagnetic waves produce electrical current in animal body, this current changes cellular function directly/indirectly; therefore, contraction/sperm discharge will be increased.

In addition, local secretion of oxytocin after exposure to electromagnetic waves has been demonstrated. This protein factor increases contractile property in seminiferous tubules [19].

**Analysis the Results of 50 Hz/4 Mt Electromagnetic Effects on Testis Weight of Mature Male Balb/C Mice Receiving Vitamin C:** According to this research, 50 Hz/4 mT electromagnetic waves

did not change significantly testis weight of mature male Balb/C mice receiving vitamin C.

**Analysis the Results of 50 Hz/4 Mt Electromagnetic Wave Effects on Body Weight of Mature Balb/C Male Mice:** According to this research, 50 Hz/4 mT electromagnetic waves reduce body weight of mature Balb/C male mice which shows significant relation.

In Louis experiments, CD-1 pregnant mice were exposed to 2/45 GHz waves from 1-17 day of pregnancy for 100 min/day, they were dissected at 18<sup>th</sup> day and fetus were analyzed in terms of abnormalities. Louis results showed that experimental fetus weight was lower than control [20]. In another study, few Spagu-Dawley rats were exposed to microwaves from 6-20 pregnancy day. Weight of test group was less that control [21]. Oconnor exposed mice to 2450 MHz and found that weight of embryos was reduced significantly because of maternal thermal stress [22]. Some researchers investigated neonate weight of female physiotherapists and concluded that their weights were less than control. They also explained thermal stress of electromagnetic waves for this weight reduction, increased temperature not only kills fetus but also delays fetus development. This phenomenon explains less weight of test neonates than control in mentioned study [23]. Dasdage et al. investigated mobile phone waves on rats, which showed decreased weight of exposed fetus [24]. In another study, invertebrate embryos exposed to low frequency electromagnetic waves lowered fertility in females and inhibited embryo development at bi cellular phase [25]. In another study, it was reported that exposure of embryonic cells to electromagnetic waves decreased cell cleavage and in harsh cases, stopped cleavage, the reason was chromosomal damage induction by electromagnetic waves [26]. It was reported that exposure of Inner Cell Mass to electromagnetic waves inhibited mitosis and pluripotency in these cells. Reason of this was free radical production in these cells [27].

In study by Rahbarian and Sadughy, it was demonstrated that development percentage of fetus exposed daily to 50 Hz and 200 Gauss electromagnetic waves was reduced significantly in comparison to fetus exposed to 10 Gauss waves [28].

Balanejad reported inhibitory effects of low frequency electromagnetic waves with 400 Gauss intensity on angiogenesis at chorioallantotic membrane of chick embryo. Balanejad believes that high intensity electromagnetic waves can reduce weight of chick embryo in early phases of growth [29]. Huuskonen claimed that exposure of pregnant wistar rats to low frequency electromagnetic waves leads to sever body weight and occur of abnormality in motor organs; in addition, high intensities lead to mortality of rats [30].

Canseven exposed embryonic cells of guinea pigs to 50 Hz with 1,2,3 T electromagnetic waves for 5 days (4 to 8 hours/day) and investigated their development. Results showed that electromagnetic radiation degenerated embryonic cells by DNA damage, influencing on membrane enzymes and changing

its permeability [31]. Cieslar studied development phases of embryonic heart cells of rats (*in vitro*) exposed to 50 Hz with 78.3 Gauss electromagnetic waves for 30 minutes. Results showed that length, diameter and size of heart in samples exposed directly to electromagnetic waves decreased significantly in comparison to control [32].

In a study, Valles showed that low frequency/high intensity electromagnetic waves could affect cell cleavage and mitotic spindle orientation; even they could inhibit cell division by damaging mitotic spindles [33]. Low frequency electromagnetic waves decrease adrenal weight and inhibit sympatho-adrenal system in hypertension rats [34]. According to Jelodar and Beizai studies, leakage waves of microwave oven reduced growth increased T<sub>3</sub>, T<sub>4</sub>, cortisol, triglyceride and HDL levels [35].

**Analysis the Results of 50 Hz/4 Mt Electromagnetic Effects on Body Weight of Mature Male Balb/C Mice Receiving Vitamin C:** According to this research, 50 Hz/4 mT electromagnetic waves increase body weight of mature Balb/C male mice which shows significant relation.

## References

1. Wood AW. How dangerous is mobile phones transmission masts and electricity pylons. Arch Dis Child. 2006; 91: 361-366.
2. Torregrossa MV. Biological and health effects on electric and magnetic fields at extremely low frequencies. Ann Ig. 2005; 17: 441-453.
3. Salzinger K. Behavioral effects of electromagnetic fields in animals. Biological effects of Electric and Magnetic fields. 1<sup>st</sup> ed. New York Academic press. 1994; 315-319.
4. Jafari-pour M, Sharafi M. Physic for radiography. The unit of Jahad daneshgahi Medical University of Iran. 1988; 89-94.
5. Polk CE. Biological effects of electromagnetic fields. 2<sup>nd</sup> ed. Boca Raton IL, CRC. 1996; 364-370.
6. Kundi M, Hardell L, Sage C, et al. Electromagnetic fields and the precautionary principle. Environ Health Perspect. 2009; 117: A484-A485.
7. Blackman CF, Benane SG, Rabinowitz JR, et al. A role for the magnetic field in the radiation induced efflux of calcium ions from brain *in vitro*. Bio electromagnetics. 1985; 8: 215-227.
8. Davanipour Z, Sobel E, Bowman JD, et al. Amyotrophic lateral sclerosis and occupational exposure to electromagnetic fields. Bio electromagnetics. 1997; 18: 25-35.
9. Vahdati A, Zandi M, Hasani F. Evaluation of electromagnetic waves from telecommunication tower.
10. Khurana VG, Teo C, Kundi M, et al. Cell phones and brain tumors a review including the long-term epidemiologic data. Surg neurol. 2009; 72: 205-214.
11. Litvak E, Foster KR, Repacholi MH. Health and safety implications of exposure to electromagnetic fields in the frequency range 300 Hz to 10 MHz. Bio electromagnetics. 2002; 23: 68-82.

12. Albert PW. Otolitic medicine and surgery. USA Churchill. 1998; 1374-1376.
13. Fritze K, Wiessner C, Kuster N, et al. Effect of global system for mobile communication microwave exposure on the genomic response of the rat brain. *Neuroscience*. 1997; 81: 627-639.
14. Reese J, Jostes R, Frazier M. Exposure of mammalian cells to 60-Hz magnetic or electric fields Analysis for DNA single-strand breaks. *Bio electromagnetics*. 1988; 9: 237-247.
15. Barnes FS. Some engineering models for interactions of electric and magnetic fields with biological systems. *Bio electromagnetics*. 1992; 53: 67-85.
16. Floderous B, Persson T, Stenlund C. Magnetic fields exposure in the workplace reference distribution and exposure in occupational groups. *Int J occup Environ Health*. 1996; 2: 226-238.
17. Savitz DA, Loomis DP. Magnetic field exposure in relation to leukaemia and brain cancer mortality among electric utility workers *Am J Epidemiol*. 1995; 141: 123-134.
18. Kaszuba-Zwoinska J, Ziomber A, Gil K, et al. Pulsating electromagnetic field induces apoptosis of rat's bowel. *Cajal's cells. Folia Med Cracor*. 2005; 46: 87-95.
19. Mezei G, Kheifets L. Selection bias and its implications for case control studies a case study of magnetic field exposure and childhood Leukemia. *Int J Epidemiol*. 2006; 35: 397-406.
20. Akdag MZ, Dasdag S, Aksen F, et al. Effect of ELF magnetic fields on lipid peroxidation sperm count p53, and trace elements. *Med Sci Monit*. 2006; 12: 366-371.
21. Christ A, Samaras T, Klingenberg A, et al. Characterization of the electromagnetic near field absorption in layered Biological tissue in the frequency range from 30 MHz to 6000 MHz. *Phys Med Biol*. 2006; 51: 4951-4965.
22. Bracken MB, Belanger K, Hellenbrank K, et al. Exposure to electromagnetic fields during pregnancy with emphasis on electrically heated beds Association with birthweight and intrauterine growth retardation. *Epidemiology*. 1995; 6: 263-270.
23. Mescher, Anthony. *Junqueira's Basic Histology*. 2011.
24. Zamiri, Mohammad Javad. *Domestic Animal Physiology*. Rasht Iran. 2000.
25. Guyton, Arthur Hall, John Edvard. *Guyton & Hall Pocket Companion to Textbook of Medical Physiology*. 2011.
26. Lieberman, Shari Pauling Bruning, Nancy. *The Real Vitamin and Mineral Book*. 2007.
27. Bordovsky SC, McCarty CA, Snibson G, et al. Management of alkali burns an 11-year retrospective review. *Ophthalmology*. 2000; 107: 1829-1835.
28. Albert PW. Otolitic medicine and surgery. USA Churchill. 1998; 1374-1376.
29. Lawrence WT, Wilmore CH. Acute wound care. *American college of surgeons. New York web MD*. 2002; 136.
30. Gandhi O, Sedigh k, Beck G. Distribution of electromagnetic energy deposition in models of man with frequencies near resonance. *Biological effects of electromagnetic waves. Selected papers of the USNGIURSI Annual Meeting. Boulder Colorado. October 20-23* *Open of Health Educ And Welfare Publ. FDA II*. 1976; 44-67.
31. Goodman R. Electromagnetic fields and cells. *J cell Biochem*. 1994; 54: 281-288.
32. Lyle DB, Fuchs TA, Casamento JP, et al. Intracellular calcium signaling by Jurkat T-Lymphocytes exposed to a 60 Hz magnetic field. *Bio electromagnetics*. 1997; 18: 439-445.
33. Parivar K, Golestanian N. The effect of variable electromagnetic field on growth on growth of bone mice with gene Balb/C. 4: 39-50.
34. Belousova TE, Kargina R. Adrenergic nerve plexuses of heart and adrenal and myocardial catecholamines of spontaneously hypertensive rats under the influence of electromagnetic irradiation in the millimeter range. *Morfologia*. 1999; 115: 16-18.
35. Mohseni Kochesfahani H, Parivar K, Golestanian N. The effect of electromagnetic fields (60 Hz) with solenoid on development of hematopoiesis system Balb/C mouse Persian. *Science Journal of Tehran University*. 2000; 26: 1-15.